

# Creating Cosmological Initial Conditions for Fun and Profit

Brian W. O'Shea

Laboratory for Computational Astrophysics

Center for Astrophysics and Space Sciences

University of California, San Diego

April 15, 2003

## **Abstract**

This is a short 'how-to' summary of one method of creating cosmological initial conditions using the Zel'Dovich approximation.

# 1 Introduction

Creating a set of initial conditions for simulations of structure formation is, on the surface, a very straightforward task. One specifies a background cosmological model, typically described as a spatially flat or open Robertson-Walker spacetime. Following that, perturbations are imposed upon this background.

The specification of background cosmology requires several inputs: The amount and nature of dark matter, the Hubble parameter  $H_0$ , and possibly the amount of baryonic matter and cosmological constant in the universe.

At the epoch of baryon-photon decoupling ( $z \sim 1100$ ), small-amplitude (“linear”) fluctuations in density are already present in all of the components of the universe (such as baryons, dark matter, photons, and neutrinos). The statistical nature of these fluctuations depends on their origin. There are two general classes of early universe models that are considered to provide reasonable mechanisms for perturbations: Topological defects (Durrer et. al 2002) and inflation (Guth 1981). Inflation predicts Gaussian fluctuations and defect models are non-Gaussian.

Gaussian fluctuations are simple since they are specified completely by a single function, the power spectrum  $P(k)$ . In Gaussian models the perturbations are set down immediately (possibly in the pre-inflationary epoch) and evolve in a straightforward manner. In real space, the probability distribution of density fluctuations is a multidimensional Gaussian, and it is very easy to sample a Gaussian random field by sampling its Fourier components on a Cartesian lattice, which is the technique that will be discussed in this paper. For more information on other methods, see Bertschinger (1998).

Non-Gaussian models are much more complicated to sample. Not only do they require more initial information than a simple power spectrum, they also are more computational effort. Typically, topological defects induce matter density fluctuations from the time of their creation in the early universe to the present day, and the dynamics of their formation and evolution are relativistic and nonlinear. For more information on creating initial conditions of topological defects, see Bertschinger (1998) or Durrer et. al (2002).

## 2 Creating Gaussian Random Fields

The creation of cosmological initial conditions using a Gaussian random field is relatively straightforward. Given a power spectrum  $P(k)$ , the linear den-

sity fluctuation field is calculated at some initial time (typically  $z \sim 100$  for high-resolution simulations). From this, dark matter particle positions and velocities are determined, along with baryon density and velocity fields. These steps are described further in Sections 2.1 and 2.2.

## 2.1 The Linear Density Fluctuation Field

The first step towards creating a gaussian density field is to specify a power spectrum. The power spectrum of the fractional density fluctuations at the redshift  $z = z_{eq}$  when the energy density in matter is equal to that in radiation, can be related to the primordial power spectrum by  $P(k, z_{eq}) = T^2(k) \times P_p(k)$ , where  $T(k)$  is the matter transfer function as a function of wave number, which describes the processing of the initial density perturbations during the radiation dominated era (Padmanabhan 1993) and  $P_p(k)$  is the primordial matter power spectrum, which typically has a power law form, ie,  $P_p(k) \sim k^n$ , where  $n$  is the index of the primordial power spectrum. This index is equal to unity for Harrison-Zel'Dovich scale-invariant spectra, a typical model. The power spectrum at any redshift  $z$  in the matter dominated era may then be written in the form

$$\frac{k^3}{2\pi^2} P(k, z) = \left( \frac{ck}{H_0} \right)^{3+n} \delta_H^2 T^2(k) D_g^2(z) / D_g^2(0), \quad (1)$$

where the  $D_g$ 's are the linear growth factor for perturbations, which is defined in Peebles (1980). A closed-form fitting function (much more appropriate for computation) is given in Eisenstein & Hu (1999).  $\delta_H$  is a constant describing the amplitude of density fluctuation, which can be provided from observations of the CMB or from large scale structure, or can be normalized by comparing to, eg,  $\sigma_8$ , which is the rms amplitude of the mass fluctuations in the universe when smoothed using a top-hat function with characteristic radius of  $8 h^{-1}$  Mpc.

Once  $P(k)$  has been determined, we then proceed to calculate  $\delta_k$ , namely, the density fluctuations in  $k$ -space. To simplify matters, we choose a three-dimensional Cartesian grid with  $N$  grid points per dimension. Each of the grid points has a unique  $(n_x, n_y, n_z)$  identifier associated with its location along the  $(x, y, z)$  axis. We sample the power spectrum  $P(k)$  discretely at each grid location  $(n_x, n_y, n_z)$ , getting  $k$  in this manner:

$$k^2 = (n_x^2 + n_y^2 + n_z^2)dk^2 \quad (2)$$

where  $dk = 2\pi/L_{box}$ , where  $L_{box}$  is the size of the simulation box in megaparsecs.  $\delta_k$  is a complex value with random amplitude and phase in a gaussian distribution with a mean of  $P(k)$ . One method to calculate it is to generate a phase angle  $\theta$ , which is randomly selected in a uniform manner in the interval  $[0, 2\pi]$ , and an amplitude  $A$  such that

$$A = \sqrt{-\log(R) * P(k)} \quad (3)$$

where  $R$  is randomly selected in a uniform manner in the interval  $(0,1)$ .  $\delta_k$  is then

$$\delta_k = Ae^{i\theta} \quad (4)$$

We then perform a Fourier transform on the grid of values of  $\delta_k$ , which gives  $\delta_x$ , the relative density fluctuation at each spatial grid point in the simulation volume. The actual physical density is then

$$\rho_{DM}(\vec{x}) = (1 + \delta_x)\overline{\rho_{DM}} \quad (5)$$

where  $\overline{\rho_{DM}}$  is the mean dark matter density in the simulation. The perturbations in the dark matter and baryon densities are assumed to be coupled, which is true in the linear regime, so the baryon density at any position is

$$\rho_b(\vec{x}) = \frac{\Omega_b}{\Omega_{DM}}\rho_{DM}(\vec{x}) \quad (6)$$

where  $\Omega_b$  and  $\Omega_{DM}$  are the ratios of the present-day mean baryon and dark matter densities to the critical density of the universe.

## 2.2 Position and Velocity Fields

The standard approach for the dark matter is to displace equal-mass particles from a uniform Cartesian lattice using the Zel'Dovich (1970) approximation:

$$\vec{x} = \vec{q} + D(t)\psi(\vec{q}) \quad (7)$$

and

$$\vec{v} = a\frac{dD}{dt}\vec{\psi} = aHfD\vec{\psi}, \quad (8)$$

where  $\vec{q}$  labels the unperturbed lattice position,  $D(t)$  is the growth factor of the linear growing mode, and  $f = d \ln D / d \ln a$  is its logarithmic growth rate (Peebles 1980). The irrotational (curl-free) displacement field  $\vec{\psi}$  is computed by solving the linearized continuity equation,

$$\vec{\nabla} \cdot \vec{\psi} = -\frac{\delta_x}{D(t)}, \quad (9)$$

Since the equation is linearized,  $\vec{\psi}$  can be found by taking the Fourier transform of  $-i\delta_k \hat{k}/k$ . See Appendix A for all equations.

### 3 Acknowledgements

Thanks to Ed Bertschinger, who wrote the excellent review article on simulation techniques of structure formation which I blatantly borrowed from. Also thanks to my labmates, (names to be determined), for proofreading this paper.

## A Some useful equations

A useful fitting function for the linear growth function  $D_g(z)$  given as

$$D_g(z) = \frac{1}{1+z} \frac{5\Omega_m(z)}{2} \left\{ \Omega_m(z)^{4/7} - \Omega_\Lambda(z) + \left[ 1 + \frac{\Omega(z)}{2} \right] \left[ 1 + \frac{\Omega_\Lambda(z)}{70} \right] \right\}^{-1} \quad (10)$$

where

$$\Omega_m(z) = \Omega_{m,0}(1+z)^3 g^{-2}(z) \quad (11)$$

and

$$\Omega_\Lambda(z) = \Omega_{\Lambda,0} g^{-2}(z) \quad (12)$$

where

$$g(z) = \Omega_{m,0}(1+z)^3 + (1 - \Omega_{m,0} - \Omega_{\Lambda,0})(1+z)^2 + \Omega_{\Lambda,0} \quad (13)$$

$\Omega_{m,0}$  and  $\Omega_{\Lambda,0}$  are the matter and vacuum energy density with respect to the critical density,  $\rho_c = 3H_0^2/8\pi G$ , at the present epoch.

The scale factor  $a(t)$  is calculated for any epoch by solving the Friedman equation,

$$\left(\frac{\dot{a}}{a}\right)^2 + \frac{k}{a^2} = \frac{8\pi G}{3}\rho + \frac{\Lambda}{3} \quad (14)$$

which comes from the  $G_0^0$  tensor component of Einstein's equation.  $\rho$  is actually a sum of the baryonic matter, dark matter, radiation, neutrino etc. contributions to the energy density at the current epoch. These quantities do not all scale the same way with  $a$  - matter scales as  $a^{-3}$ , while radiation scales as  $a^{-4}$ . This equation can be rewritten as:

$$\left(\frac{\dot{a}}{a}\right)^2 + \frac{k}{a^2} = H_0^2 \left( \frac{\Omega_{r,0}}{a^3} + \frac{\Omega_{m,0}}{a^3} + \Omega_\Lambda \right) \quad (15)$$

In general an exact solution of this can only be obtained numerically.

## References

- [] Bertschinger, E. 1998 ARAA, 36, 599-654
- [] Durrer et al. 2002 Phys.Rept., 1-81
- [] Eisenstein, E. & Hu, W. 1999 ApJ, 511, 5
- [] Guth, 1981 Phys. Rev. D, 23, 347-356
- [] Majumdar, S. & Subrahmanyam, R. 2000 MNRAS, 312, 724
- [] Padmanabhan, T. 1993 Structure Formation In The Universe, Cambridge University Press, Cambridge, England
- [] Peebles, P. 1980 The Large Scale Structure of the Universe, Princeton University Press, Princeton
- [] Zel'Dovich, Y. 1970 Astron. Astrophys., 5, 84-89